

Variational Calculation on the Helium Atom Using a Hydrogenic Trial Wave Function

Frank Rioux
Chemistry Department
CSB|SJU

Gaussian Trial Wave Function:
$$\Psi(r, \beta) := \sqrt{\frac{\beta^3}{\pi}} \cdot \exp(-\beta \cdot r)$$

Demonstrate the wave function is normalized.
$$\int_0^{\infty} \Psi(r, \beta)^2 \cdot 4 \cdot \pi \cdot r^2 dr \left| \begin{array}{l} \text{assume, } \beta > 0 \\ \text{simplify} \end{array} \right. \rightarrow 1$$

The terms contributing to the total electronic energy of the helium atom are the kinetic energy of each electron, each electron's interaction with the nucleus, and the interaction of the electrons with each other.

Calculate kinetic energy:
$$2 \cdot \int_0^{\infty} \Psi(r, \beta) \cdot \left[-\frac{1}{2 \cdot r} \cdot \frac{d^2}{dr^2} (r \cdot \Psi(r, \beta)) \right] \cdot 4 \cdot \pi \cdot r^2 dr \left| \begin{array}{l} \text{assume, } \beta > 0 \\ \text{simplify} \end{array} \right. \rightarrow \beta^2$$

Calculate electron-nucleus potential energy:

$$2 \cdot \int_0^{\infty} \Psi(r, \beta) \cdot \frac{-2}{r} \cdot \Psi(r, \beta) \cdot 4 \cdot \pi \cdot r^2 dr \left| \begin{array}{l} \text{assume, } \beta > 0 \\ \text{simplify} \end{array} \right. \rightarrow (-4) \cdot \beta$$

Calculation of electron-electron potential energy:

a. Calculate the electric potential of one of the electrons in the presence of the other:

$$\frac{1}{r} \cdot \int_0^r \Psi(x, \beta)^2 \cdot 4 \cdot \pi \cdot x^2 dx + \int_r^{\infty} \frac{\Psi(x, \beta)^2 \cdot 4 \cdot \pi \cdot x^2}{x} dx \left| \begin{array}{l} \text{assume, } \beta > 0 \\ \text{simplify} \end{array} \right. \rightarrow \frac{-[\beta \cdot r \cdot e^{(-2) \cdot r \cdot \beta} + e^{(-2) \cdot r \cdot \beta} - 1]}{r}$$

b. Calculate the electron-electron potential energy using result of part a:

$$\int_0^{\infty} \Psi(r, \beta)^2 \cdot \left[\frac{-[\beta \cdot r \cdot e^{(-2) \cdot r \cdot \beta} + e^{(-2) \cdot r \cdot \beta} - 1]}{r} \right] \cdot 4 \cdot \pi \cdot r^2 dr \left| \begin{array}{l} \text{assume, } \beta > 0 \\ \text{simplify} \end{array} \right. \rightarrow \frac{5}{8} \cdot \beta$$

Write the equation for the total electronic energy in terms of the variational parameter β and minimize the energy with respect to β :

$$E(\beta) := \beta^2 - 4\beta + \frac{5}{8}\beta \quad \beta := \frac{d}{d\beta}E(\beta) = 0 \text{ solve, } \beta \rightarrow \frac{27}{16} \quad E(\beta) = -2.848$$

Compare the variational calculation to the Hartree-Fock limit: $E_{\text{HF}} := -2.8617$

$$\frac{E_{\text{HF}} - E(\beta)}{E_{\text{HF}}} = 0.491 \%$$

Compare optimized trial wave function with the Hartree-Fock wave function (see McQuarrie and Simon, page 283) by plotting the radial distribution functions.

$$\Phi(r) := .75738 \cdot \exp(-1.430 \cdot r) + .43658 \cdot \exp(-2.4415 \cdot r) + .17295 \cdot \exp(-4.0996 \cdot r) - .02730 \cdot \exp(-6.4843 \cdot r) + .06675 \cdot \exp(-7.978 \cdot r)$$

