

The Difference Between Fermions and Bosons

$$n_1 := 1 \quad n_2 := 2 \quad \Psi(x) := \sqrt{2} \cdot \sin(n_1 \cdot \pi \cdot x) \quad \Phi(x) := \sqrt{2} \cdot \sin(n_2 \cdot \pi \cdot x)$$

Calculate the average separation, $|x_1 - x_2|$, for two fermions and two bosons in a 1D box of unit length.

Fermions have antisymmetric wave functions:


$$\Psi_f(x_1, x_2) := \frac{\Psi(x_1) \cdot \Phi(x_2) - \Psi(x_2) \cdot \Phi(x_1)}{\sqrt{2}}$$

The average particle separation for indistinguishable fermions:

$$\int_0^1 \int_0^1 \Psi_f(x_1, x_2) \cdot |x_1 - x_2| \cdot \Psi_f(x_1, x_2) dx_1 dx_2 = 0.383$$

The particles are correlated so as to keep them apart.

$$N := 40 \quad i := 0..N \quad x_{1_i} := \frac{i}{N} \quad j := 0..N \quad x_{2_j} := \frac{j}{N} \quad \Psi_{f_{i,j}} := \Psi_f(x_{1_i}, x_{2_j})^2$$



Bosons have symmetric wave functions:

$$\Psi_b(x_1, x_2) := \frac{\Psi(x_1) \cdot \Phi(x_2) + \Psi(x_2) \cdot \Phi(x_1)}{\sqrt{2}}$$


The average particle separation for indistinguishable bosons:

$$\int_0^1 \int_0^1 \Psi_b(x_1, x_2) \cdot |x_1 - x_2| \cdot \Psi_b(x_1, x_2) dx_1 dx_2 = 0.157$$


The particles are correlated so as to bring them closer together.

$$N := 40 \quad i := 0..N \quad x_{1_i} := \frac{i}{N} \quad j := 0..N \quad x_{2_j} := \frac{j}{N} \quad \Psi_{b_{i,j}} := \Psi_b(x_{1_i}, x_{2_j})^2$$

x_1



Ψ_b



x_2

Ψ_b

All fundamental particles (electrons, neutrons, protons, photons, etc.) are either bosons or fermions. Composite entities such as the elements also fall into these two categories. The fundamental distinction is spin: bosons have integer spin (0, 1, 2, ...) while fermions have half-integer spin (1/2, 3/2, ...).

The dramatic difference in behavior between bosons and fermions has led to a sociology of fundamental particles. Bosons are social and gregarious, while fermions are antisocial and aloof.